



The Physics potential of LHCb: CP violation and rare decays in the beauty sector at LHCb

R. Le Gac

► To cite this version:

R. Le Gac. The Physics potential of LHCb: CP violation and rare decays in the beauty sector at LHCb. Les Rencontres de physique de la Vallée d'Aoste : Results and Perspectives in Particle Physics, Feb 2000, La Thuile, Italy. pp.1-21. in2p3-00011371

HAL Id: in2p3-00011371

<https://hal.in2p3.fr/in2p3-00011371>

Submitted on 22 Mar 2002

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The Physics Potential of LHCb:

CP violation and rare decays in the beauty sector at LHC

R. Le Gac[†]
on behalf of the LHCb Collaboration

1. Motivation
2. Methods
3. LHCb detector
4. Physics performances
5. Status and Conclusions


[†]. Centre de Physique des Particules de Marseille, IN2P3–CNRS



MOTIVATION

- Standard Model accommodates CP violation through the CKM matrix. This mechanism has to be validated.

Unitarity property of the matrix \rightarrow Unitarity Triangles (UT):



- Currently the UT is weakly constrained (ϵ , $|V_{ub}|$, $|V_{cb}|$, Δm_d)
- Will be improved in ~2005:
 - Δm_s ? (HERA-b, CDF/D0)
- CP violation in the B_d system:
 - β (BaBar, Belle, CDF/D0, HERA-b)
 - α ? (BaBar, Belle, HERA-b)
- Significant theoretical uncertainties on the length of the sides

- LHCb will measure:

- CP violation in the B_s system
- all phases: β , γ , $\delta\gamma$, $(\beta+\gamma)$,...
- one side: Δm_s

\rightarrow Over-constrain the unitarity triangles.

Understand CP violation in the framework of the Standard Model.

MOTIVATION ...

- **2nd motivation is related to the evolution of the Universe.**
The disappearance of the antimatter component requires CP violation.
But, the CP violation from the Standard Model is too small.
 - ➔ **Additional sources of CP violation are needed.**
They might find their origin in supersymmetric models
- **LHCb is a promising place to study new physics.**
Complementary approach to ATLAS and CMS.
- **LHCb will measure:**
 - The phase β, γ using different decay modes.
 - Branching ratio of rare decays such as: $B_s \rightarrow \mu^+ \mu^-$, $B_{d,s} \rightarrow K^{*0} \mu^+ \mu^-$, ...

These measurements couple with the unitarity checks of the CKM matrix will allow to pin down the details of new physics.

2.

Experimental methods
to determine
 β , γ , $(\beta+\gamma)$, $\delta\gamma$,...

TIME DEPENDENT DECAY RATE ASYMMETRY

- $$A_{CP}(\tau) = \frac{\Gamma(B_{d,s}^0 \rightarrow f_{CP})(\tau) - \Gamma(\bar{B}_{d,s}^0 \rightarrow f_{CP})(\tau)}{\Gamma(B_{d,s}^0 \rightarrow f_{CP})(\tau) + \Gamma(\bar{B}_{d,s}^0 \rightarrow f_{CP})(\tau)} \neq 0 \Leftrightarrow \begin{cases} \text{CP Violation} \\ \text{CP}|B^0\rangle = |\bar{B}^0\rangle \\ \text{CP}|f_{CP}\rangle = \pm|f_{CP}\rangle \end{cases}$$

- $$A_{CP}(\tau) \equiv \begin{array}{c} \text{B}^0 \\ \text{B}^0 \end{array} \begin{array}{c} \xrightarrow{\text{orange}} \\ \xrightarrow{\text{green}} \end{array} \begin{array}{c} f_{CP} \\ \bar{B}^0 \end{array} \equiv \pm \sin[\underbrace{2(\phi_M + \phi_D)}_{\text{orange}}] \sin(\Delta m \tau)^\dagger$$

		Visible BR
$B_d^0 \rightarrow J/\psi K_s$	2β	3.6×10^{-5}
$B_d^0 \rightarrow \pi^+ \pi^-$	$2\beta + 2\gamma$	0.7×10^{-5}
$B_s^0 \rightarrow J/\psi \phi$	$2\delta\gamma$	5.4×10^{-5}
	$\left \frac{P}{T} \right , \delta_{P/T}$	
	$\left \frac{A(\text{CP}=+1)}{A(\text{CP}=-1)} \right , \delta_1, \Delta\Gamma_s$	

†. When more than one amplitude contributes to the decay: $A_{CP}(\tau) = A_{f_{CP}}^{dir} \cos(\Delta m \tau) + A_{f_{CP}}^{mix} \sin(\Delta m \tau)$

TIME DEPENDENT DECAY RATE ASYMMETRY...

- When the final state is not a CP eigenstate:

			Visible BR
$B_S^0 \rightarrow D_S^\pm K^\mp$	$-2\delta\gamma + \gamma$	$\left \frac{T_1}{T_2} \right , \delta_{T_1/T_2}$	0.9×10^{-5}
$B_d^0 \rightarrow D^{*\pm} \pi^\mp$	$2\beta + \gamma$	$\left \frac{T'_1}{T'_2} \right , \delta_{T'_1/T'_2}$	6.7×10^{-5}
$B_S^0 \rightarrow D_S^- \pi^+$		$\Delta m_S, \Delta \Gamma_S$	12×10^{-5}

OTHER METHODS...

- Dalitz plot analysis:

				Visible BR
$B_d^0 \rightarrow \rho \pi \rightarrow \pi^+ \pi^- \pi^0$	$\beta + \gamma$	$T^{+-}, T^{00}, P^{+-}, P^{-+}$	5.5×10^{-5}	


- Branching ratio measurements:

				Visible BR
$B_d^0 \rightarrow \begin{cases} \bar{D}^0 K^{*0} \\ D^0 K^{*0} + \text{c.c.} \\ D_1 K^{*0} \end{cases}$	γ	δ_4	$10^{-8} - 10^{-7}$	

- New strategies are proposed to measure γ .
They are under evaluation.

3.

LHCb detector...



LHCb DETECTOR PERFORMANCES

- Particle identification [RICH]:

- K/π separation $> 3\sigma$ in the momentum range of the experiment

- Resolutions [Vertex detector, Tracking chambers]:

- Momentum $\approx 0.3\%$
- Impact parameter $\approx 40 \mu\text{m}$
- Decay-time $\approx 0.03 \tau_B$ (43 fs)

- Tagging:

- Efficiency (K, e, μ) $\approx 40\%$
- Dilution factor $\approx 40\%$

RUNNING CONDITIONS

Low Luminosity: $2 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$

- **Tunable locally although ATLAS and CMS work at high luminosity**
 - Ensures long physics programme
 - Guarantees physics results from the beginning of LHC
- **Clean event:**
 - ~ 0.3 interactions/BC
- **Highest statistics of B's ever collected:**
 - 4.5×10^{11} $B_d + \bar{B}_d$ produce per year
 - 1.3×10^{11} $B_s + \bar{B}_s$ produce per year



4.

Physics performances...

$$B_d \rightarrow \pi^+ \pi^- (\beta + \gamma)$$

Importance of particle identification


Branching ratio: $B_d \rightarrow \pi^+ \pi^- = 0.7 \times 10^{-5}$, $\rightarrow K^\pm \pi^\mp = 1.5 \times 10^{-5}$
 $B_s \rightarrow K^+ K^- = 1.5 \times 10^{-5}$, $\rightarrow K^\pm \pi^\mp = 0.7 \times 10^{-5}$



$$B_s \rightarrow D_s K (\gamma - 2\delta\gamma)$$


Major background: $B_s \rightarrow D_s \pi$ (No CP violation)

Importance of particle identification and mass resolution



$$B_d \rightarrow \rho\pi \rightarrow \pi^+\pi^-\pi^0 \quad (\beta+\gamma), |P|, |T|, \dots$$

Importance of π^0 reconstruction




Events per year:

$B_d \rightarrow \rho^+\pi^-$	~ 1000	(Br 4.4×10^{-5})
$B_d \rightarrow \rho^-\pi^+$	~ 200	(Br 1.0×10^{-5})
$B_d \rightarrow \rho^0\pi^0$	~ 100	(Br 0.1×10^{-5})

$$B_s \rightarrow D_s \pi (\Delta M_s, \Delta \Gamma_s)$$

Importance of decay-time resolution



➔ Measurement of Δm_s with a significance > 5 , up to 48 ps^{-1} ($\chi_s=75$)
Standard Model expectation: $14.3\text{--}26 \text{ ps}^{-1}$

SUMMARY OF THE PERFORMANCES

Sensitivity per year of data taking:

LHCb Lum=2×10 ³²			
$B_d^0 \rightarrow J/\psi K_S$	$\sigma(\sin 2\beta)$	0.02	when $\beta = 20^\circ$
$B_d^0 \rightarrow \pi^+ \pi^-$	$\sigma(A^{\text{dir}})$	0.07	
	$\sigma(A^{\text{mix}})$	0.09	
	$\sigma(\beta + \gamma)$	$\sim 2^\circ$	If $ P/T = 0.20 \pm 0.02$ when $\alpha = 90^\circ$ and $\delta = 30^\circ$
$B_d^0 \rightarrow \rho \pi$	$\sigma(\beta + \gamma)$	$2^\circ - 5^\circ$	Preliminary: Backg. not included in the fit Depends on the value of $\beta + \gamma$
$B_d^0 \rightarrow D^{*\pm} \pi^\mp$	$\sigma(2\beta + \gamma)$	$\sim 9^\circ$	when $ 2\beta + \gamma < 25^\circ$
$B_S^0 \rightarrow J/\psi \phi$	$\sigma(\delta\gamma)$	$0.6^\circ - 1.4^\circ$	depends on the value of Δm_s SM expectation: $\sim 1^\circ$
$B_S^0 \rightarrow D_S^\pm K^\mp$	$\sigma(\gamma - 2\delta\gamma)$	$6^\circ - 13^\circ$	Depends on the value of: • $\gamma - 2\delta\gamma$ • strong phase shift difference
$B_d^0 \rightarrow D^0 K^{*0}$	$\sigma(\gamma)$	$4^\circ - 19^\circ$	Depends on the value of: • γ • strong phase shift difference

$B_S^0 \rightarrow D_S^\pm \pi^\mp$	x_s reach	<75	Significance > 5
$B_S^0 \rightarrow \mu^+ \mu^-$	S/\sqrt{B}	~ 6	SM expectation BR = 3.5×10^{-9}

$B_d^0 \rightarrow K^{*0} \gamma$	$ V_{ts} $	26×10^3 evts.	
-----------------------------------	------------	------------------------	--

5.


Status and conclusions...



Brazil



Finland



France



Germany



Italy




Poland



PRC



Netherlands



Romania




Russia



Spain



The *LHCb* Experiment



USA




Ukraine



UK




Switzerland



STATUS AND PLANS

- Technical Proposal approved in September 98

- Preparing Technical Design Reports:



- Construction: 2001–2003

- Ready for the first collision of the LHC

CONCLUSIONS

- LHCb is an opportunity to understand the CP violation in the framework of the Standard Model and BEYOND.
- LHCb fully exploit the physics from the beginning of the LHC